

(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 829 463 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

18.03.1998 Bulletin 1998/12

(51) Int Cl.⁶: C07C 33/12, C11B 9/00

(21) Application number: 97307074.1

(22) Date of filing: 11.09.1997

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

Designated Extension States:

AL LT LV RO SI

(30) Priority: 17.09.1996 JP 244842/96

16.04.1997 JP 98760/97

(71) Applicant: Takasago International Corporation
Tokyo 108 (JP)

(72) Inventors:

- Aida, Takashi, c/o Takasago Int. Corp.
Hiratsuka-shi, Kanagawa (JP)

• Harada, Makoto, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

• Iwai, Hisao, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

• Amano, Akira, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

• Yamasaki, Tetsuro, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

• Yamamoto, Takeshi

Minato-ku, Tokyo (JP)

(74) Representative: Bubbs, Antony John Allen et al
GEE & CO.


Chancery House

Chancery Lane

London WC2A 1QU (GB)

(54) (E)-(R)-2-Alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol, process for preparing the same, and use thereof in perfume compositions

(57) An (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol represented by formula (I):



(I)

wherein R represents an alkyl group having 1 to 3 carbon atoms,

and a process for preparing the same comprising hydrogenating a corresponding (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol in the presence of a ruthenium-phosphine complex as a catalyst, a base comprising an alkali metal or an alkaline earth metal, and an amine. The compound (I) has an excellent sandalwood oil odor.

EP 0 829 463 A2

Description

This invention relates to an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol which is a novel optically active compound, a process for preparing the same, and a perfume composition containing the same.

An (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol has a fine sandalwood-like odor, and a perfume composition containing it is useful in various cosmetics, fragrances, toiletries, and other hygienic goods.

Sandalwood oil is an essential oil obtained from sandalwood tree occurring in the east of India and is highly valued as a perfume component. However, production of sandalwood oil is being decreased because, felling of the sandalwood tree is restricted from the standpoint of environmental and resources conservation pressures. Therefore synthetic sandalwood oil fragrance materials have been developed for substitution.

It is known that the main components responsible for the odor of natural sandalwood oil are α -santalol and β -santalol. Santalol is difficult to synthesize in quantities because of its chemical structure. In recent years, substances different from santalol in chemical structure but very similar in odor to santalol have been synthesized.

Among such substances are 2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol derivatives (see JP-B-56-361676 and JP-A-55-139330, the term "JP-B" as used herein means an "examined Japanese patent publication", and the term "JP-A" as used herein means an "unexamined published Japanese patent application"). 2-Methyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol is available from Taiyo Koryo K.K. under the trade name of "Santalinol", and 2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol from International Flavor and Fragrance (hereinafter abbreviated as IFF) under the registered trademark "Bacdanol".

There is found no report on synthesis of an optically active 2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol.

R. Reinhardt, et al. have reported in *J. Chromatog. A*, Vol. 697, No. 7, pp. 475-484 (1995) that the resolution of 2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol was not successful by gas chromatography using a chiral column.

Thus, there is no report on the isolation or synthesis of an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol, i.e., an optically active compound of a 2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol in which the trimethylcyclopentene ring and the hydroxymethyl group are in a trans configuration.

Several reports are found with respect to the relation between the configurations of the side chain of fragrance substances having a trimethylcyclopentene ring and the odor.

For example, (1) concerning the relation between odor and the absolute configuration of the asymmetric carbon atom on the 1-position of the cyclopentene ring in compounds having a (2,2,3-trimethyl-3-cyclopenten-1-yl)alkanol skeleton, U.S. Patent 4,052,341 and U.S. Patent 4,610,813 have reported that the (R)-isomer and the (S)-isomer have the same odor as the racemate, that is, the absolute configuration has no influence on odor.

(2) As for 3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-4-penten-2-ol, U.S. Patent 4,696,766 has disclosed that the relationship of the odor and relative configuration of 2 and 3-positions that a mixture of anti-diastereomers is 500 times that of a syn-form diastereomer mixture, but has no mention of the relation between the absolute configuration of the 1-position of the trimethylcyclopentene ring and the odor.

(3) Regarding the four optical isomers and diastereomer mixtures of (E)-3,3-dimethyl-5-(2,2,3-trimethylcyclopent-3-en-1-yl)-4-penten-2-ol, U.S. Patent 5,512,543 has described that the (1R)-isomer (mixture of diastereomers in which the 2-position is racemic) has a milky and strong sandalwood-like odor; the (1S)-isomer (mixture of diastereomers in which the 2-position is racemic) is drier than the (1R)-isomer and has a sandal cedar note; the (1S,2S)-isomer has a milky and strong sandalwood oil odor with a faint animal note; the (1S,2R)-isomer has a weak and flat sandalwood-like odor; the (1R,2S)-isomer has a more pleasant odor than the (1R,2R)-isomer and bears the odor of the (1R)-isomer (diastereomer mixture in which the 2-position is racemic); and a (2S)-isomer (mixture of diastereomers in which the 1-position is racemic) has a pleasant odor.


SUMMARY OF THE INVENTION

An object of the present invention is to provide an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol which is different from known racemic mixtures and has an excellent odor and a perfume composition containing the same.

As a result of extensive study, the inventors of the present invention have found that an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol can easily be obtained by hydrogenating an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al (hereinafter sometimes referred to as the (E-R) compound) in the presence of a base, an amine, and a ruthenium-phosphine complex catalyst. They have synthesized an (E)-(S)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol, which is an antipode of the (E-R)-compound, and have these optical isomers and their racemic mixture evaluated in terms of odor by specialized panelists. As a result, they found that the configuration of the 1-position of the cyclopentene ring is greatly influential on the odor.


That is, the novel (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol was found to have markedly excellent odor compared with its diastereomer or racemic mixture and to be very useful as a perfume material having a sandalwood oil odor. The present invention has been completed based on this finding.

The present invention provides an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol represented by formula (I):



(I)

wherein R represents a straight-chain or branched alkyl group having 1 to 3 carbon atoms; a process for preparing the compound of formula (I) comprising hydrogenating an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al represented by formula (II):



(II)

wherein R is as defined above, in the presence of a ruthenium-phosphine complex as a catalyst, a base comprising an alkali metal or an alkaline earth metal, and an amine; and a perfume composition containing the compound of formula (I).


DETAILED DESCRIPTION OF THE INVENTION

The alkyl group of the (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol of formula (I) includes a methyl group, an ethyl group, a propyl group, and an isopropyl group. (E)-(R)-2-Methyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol gives off a fresh sandalwood oil odor, and (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol has a strong, clean and bright odor. The latter compound is preferred.

The term "(E)-isomer" as used herein denotes a compound in which the trimethylcyclopentene ring and the formyl group or hydroxymethyl group are in a trans configuration.

The term "optically active" is used herein for a substance having an optical purity (enantiomer excess: % e.e.) of at least 50% e.e. The optical purity of an (R)-isomer is obtained by subtracting an (S)-isomer content (%) from an (R)-isomer content (%); and that of an (S)-isomer is calculated by subtracting an (R)-isomer content (%) from an (S)-isomer content (%).

A 2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol can be synthesized through known routes. The following process is known as one of them.



wherein R is as defined above.

That is, commercially available α -pinene (III) is epoxidized, and the resulting α -pinene oxide (IV) is subjected to transformation reaction using a Lewis acid, such as zinc bromide, to obtain camphorenic aldehyde (V) (see J.B. Lewis, et al., *J. Org. Chem.*, Vol. 30, pp. 4271-4275 (1965)). An Aldol reaction between camphorenic aldehyde (V) and an aldehyde gives a 2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al (VI) (see German Patent 68936). While the compound (VI) as obtained is usually a mixture of an (E)-isomer and a (Z)-isomer, when the alkyl group on the 2-position is as small as specified in the present invention, the steric hindrance is small so that a thermodynamically stable (E)-isomer is produced preferentially.

There have been many reports on the hydrogenation of the formyl compound (VI) into an (E)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (VII), but the known hydrogenation techniques have their several disadvantages. For example, catalytic hydrogenation using a solid catalyst, such as a Raney nickel catalyst, a copper-chromium catalyst, and a Raney copper catalyst, has the disadvantage of being accompanied by hydrogenation of the double bond in competition with reduction of the aldehyde group, resulting in a low purity of the product. Chemical reduction using lithium aluminum hydride or sodium borohydride or Meerwein-Ponndorf reduction using an aluminum alkoxide are superior in the regio-selectivity to an aldehyde group. However, the process of using lithium aluminum hydride is unsuitable for mass production because of the expensiveness and high danger of ignition of lithium aluminum hydride. The processes using sodium borohydride or an aluminum alkoxide require many steps for isolating a pure product, and the reagents used are expensive, too. Therefore, the yield attained is low, and the cost of production is high.

In contrast, the hydrogenation process according to the present invention which is carried out in the presence of a ruthenium-phosphine complex as a catalyst and a base comprising an alkali metal or an alkaline earth metal and an amine has the advantages of simple steps, inexpensive reagents, and reduction selectivity to the ketone moiety, providing the compound (I) in high yield.

The compound (II) which can be used in the present invention as a starting material is prepared from commercially available (1S,5S)- α -pinene having formula (VIII) in accordance with the above-described known process.




(VIII)

According to the process of the invention the compound (I) can be obtained by retention of the chirality of (1S,5S)- α -pinene (VIII). The optical purity of the compound (I) is dependent on that of the compound (II), and the optical purity of the compound (II) is dependent on that of (1S,5S)- α -pinene. The inventors have experimentally confirmed that the optical purity of the compound (I) is in agreement with that of the α -pinene used as a starting material.

More specifically, the inventors synthesized (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol


(IX) and (E)-(S)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (XI) from (1S,5S)- α -pinene and (1R,5R)- α -pinene, respectively. The resulting compounds (IX) and (XI) were esterified with (R)-2-methoxy-2-phenyl-3,3,3-trifluoropropionic acid (hereinafter abbreviated as MTPA) to obtain esters of the respective diastereomers.



The $^1\text{H-NMR}$ spectrum of a mixture of the (R)-(R)-diastereomer (X) and the (S)-(R)-diastereomer (XII) exhibits 8 peaks in two sets of a quartet having different coupling constants assigned to the methylene group on 1-position. Each of the compounds (X) and (XII) obtained by the process of the invention shows only 4 peaks, proving to be a single diastereomer. It has now been confirmed that the stereoisomerism of the starting α -pinene remains in the final product.

Accordingly, in order to obtain the compound (I) having an optical purity of 50%e.e. or higher, the starting (1S,5S)- α -pinene should have an optical purity of 50%e.e. or higher.

Synthesis of the compound (II) from (1S,5S)- α -pinene according to a known process is accompanied with by-production of a small amount of a (Z)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (XIII). The compound (II) is led to the compound (I) by hydrogenation, while the compound (XIII) is led to a corresponding (Z)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (XIV) as illustrated below. In other words, the process of the present invention guarantees that all stereocenters are retained throughout the reactions involved.



wherein R is as defined above.

As mentioned above, when the process of the invention is followed, there is obtained a mixture consisting of the compound (I) ((E)-R-compound) and a small proportion of the (Z)-(R)-compound (XIV). The inventors have isolated each compound from the mixture and determined their structures by two-dimensional NMR.

The (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (I) was synthesized for the first time by the

inventors.


The ruthenium-phosphine complex which is used as a catalyst for hydrogenating the compound (II) is a compound composed of ruthenium with a phosphine compound, preferably an organic phosphine compound, coordinated as a ligand. The complex may have an auxiliary ligand. It may be either a mononuclear complex or a polynuclear complex.

Commercially available ruthenium-phosphine complexes can be utilized as such, or a complex may be prepared on use. The complex can be prepared by adding 1 to 4 equivalents of a ligand compound to ruthenium. Alternatively, the complex may be prepared *in situ* by separately adding a ruthenium salt or a ruthenium complex and a ligand to the hydrogenation reaction system. Complexes having the ligand in excess, those to which triethylamine or a Lewis acid has been added, or those activated by reduction may also be used.

The organic phosphine compound coordinating to ruthenium may be either monodentate or polydentate. Monodentate ligands include those represented by formula (XV):



wherein R^1 , R^2 , and R^3 , which may be the same or different, each represent a substituted or unsubstituted alkyl group, a substituted or unsubstituted aralkyl group or a substituted or unsubstituted aryl group; and bidentate ligands include those represented by formula (XVI):



wherein R^4 , R^5 , R^6 , and R^7 , which may be the same or different, each represent a substituted or unsubstituted alkyl group, a substituted or unsubstituted aralkyl group or a substituted or unsubstituted aryl group; and A^1 represents a substituted or unsubstituted alkylene group, $-A^2-Ar-Ar-A^2-$ or $-Ar-Ar-$, wherein A^2 represents a substituted or unsubstituted alkylene group; and $-Ar-Ar-$ represents a 1,1'-biphenylene group having a bond at the 2,2'-positions, a 1,1'-binaphthylene group having a bond at the 2,2'-positions, or 5,5',6,6',7,7',8,8'-octahydro-1,1'-binaphthylene group having a bond at the 2,2'-positions, in which the biphenylene group may be substituted with a methyl group, a methoxy group or a dialkylamino group, and the binaphthylene group may be substituted with an alkali sulfonate group.

The substituted or unsubstituted alkyl group as represented by R^1 , R^2 , R^3 , R^4 , R^5 , R^6 or R^7 includes a straight-chain, branched or cyclic alkyl group which may have one or more substituents, such as a halogen atom and an alkoxy group, preferably a straight-chain or branched alkyl group having 1 to 10 carbon atoms and a cyclic alkyl group having 3 to 8 carbon atoms. Specific examples of the alkyl group are methyl, ethyl, butyl, octyl, and cyclohexyl groups.

The substituted or unsubstituted aralkyl group as represented by R^1 , R^2 , R^3 , R^4 , R^5 , R^6 or R^7 includes an alkyl group (preferably an alkyl group having 1 to 4 carbon atoms) substituted by an aryl group (preferably a phenyl group or a naphthyl group) which may have one or more substituents, such as a halogen atom, an alkyl group, and an alkoxy group. Specific examples of the aralkyl group are benzyl, phenethyl and naphthylmethyl groups.

The substituted or unsubstituted aryl group as represented by R^1 , R^2 , R^3 , R^4 , R^5 , R^6 or R^7 includes an aryl group which may have one or more substituents, such as a halogen atom, an alkyl group, and an alkoxy group, preferably a phenyl or naphthyl group which may be substituted with a chlorine atom, a bromine atom, an iodine atom, a fluorine atom, a straight-chain or branched alkyl group having 1 to 4 carbon atoms or a straight-chain or branched alkoxy group having 1 to 4 carbon atoms. Specific examples of the aryl group include phenyl, naphthyl, p-chlorophenyl, p-bromophenyl, p-fluorophenyl, p-tolyl, p-t-butylphenyl, 3,5-dimethylphenyl, and p-methoxyphenyl groups, with phenyl and p-tolyl groups being particularly preferred.

The substituted or unsubstituted alkylene group as represented by A^1 or A^2 denotes a straight-chain or branched alkylene group which may have one or more substituents, such as a halogen atom and an alkoxy group, preferably an unsubstituted straight-chain or branched alkylene group having 1 to 5 carbon atoms. Specific examples of A^1 are an ethylene group $-(CH_2)_2-$, a trimethylene group (or 1,3-propylene group) $-(CH_2)_3-$, a tetramethylene group (or 1,4-butylene group) $-(CH_2)_4-$, and a dimethylethylene group (or 2,3-butylene group) $-(CH(CH_3)CH(CH_3)-)$. Specific examples of A^2 include a methylene group $-(CH_2)-$.

Examples of preferred ligands represented by formula (XV) are trimethylphosphine, triethylphosphine, tributylphosphine, trioctylphosphine, tricyclohexylphosphine, tribenzylphosphine, triphenylphosphine, tri(p-chlorophenyl)phosphine, tri(p-bromophenyl)phosphine, tri(p-fluorophenyl)phosphine, tri(p-tolyl)phosphine, tri(p-t-butylphenyl)phosphine, tri(3,5-dimethylphenyl)phosphine, tri(p-methoxyphenyl)phosphine, methyldiphenylphosphine, and dimethylphenylphosphine.

Of the ligands represented by formula (XVI), those in which A^1 is a substituted or unsubstituted alkylene group preferably include 1,2-bis(dimethylphosphino)ethylene, 1,3-bis(dimethylphosphino)propylene, 1,4-bis(dimethylphosphino)butylene, and 1,5-bis(dimethylphosphino)pentylene.

phino)butyl n, 1,2-bis(diphenylphosphino)ethylene (hereinafter abbreviated as DPPE), 1,3-bis(diphenylphosphino)propylene (hereinafter abbreviated as DPPP), 1,4-bis(diphenylphosphino)butylene (hereinafter abbreviated as DPPB), 1,2-bis[di(p-tolyl)phosphino]ethylene, 1,3-bis[di(p-tolyl)phosphino]propylene, 1,4-bis[di(p-tolyl)phosphino]butylene, and 2,3-bis(diphenylphosphino)butylene (hereinafter referred to as CHIRAPHOS).

Of the ligands represented by formula (XVI), those in which A¹ is -A²-Ar-Ar-A²- preferably include 2,2'-bis(diphenylphosphinomethyl)-1,1'-biphenyl and 2,2'-bis(diphenylphosphinomethyl)-1,1'-binaphthyl.

Of the ligands represented by formula (XVI), those in which A¹ is -Ar-Ar- preferably include 2,2'-bis(dicyclohexylphosphino)-6,6'-dimethyl-1,1'-biphenyl (hereinafter referred to as BICHEP), 6,6'-bis(diphenylphosphino)-2,2'-dimethyl-1,1'-biphenyl (hereinafter referred to as BIPHEMP), 6,6'-bis(diphenylphosphino)-2,2'-dimethoxy-1,1'-biphenyl, 2,2',4,4'-tetramethoxy-6,6'-bis(diphenylphosphino)-1,1'-biphenyl, 4,4'-bis(dimethylamino)-6,6'-bis(diphenylphosphino)-2,2'-dimethyl-1,1'-biphenyl, 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (hereinafter referred to as BINAP), 2,2'-bis(di-p-tolylphosphino)-1,1'-binaphthyl (hereinafter referred to as Tol-BINAP), 2,2'-bis(di-m-tolylphosphino)-1,1'-binaphthyl (hereinafter referred to as m-Tol-BINAP), 2,2'-bis(di-p-t-butylphenylphosphino)-1,1'-binaphthyl (hereinafter referred to as t-Bu-BINAP), 2,2'-bis[di-(3,5-dimethylphenyl)phosphino]-1,1'-binaphthyl (hereinafter referred to as DM-BINAP), 2,2'-bis(di-p-methoxyphenylphosphino)-1,1'-binaphthyl (hereinafter referred to as MeO-BINAP), 2,2'-bis(di-p-chlorophenylphosphino)-1,1'-binaphthyl (hereinafter referred to as Cl-BINAP), 2,2'-bis(dicyclopentylphosphino)-1,1'-binaphthyl (hereinafter referred to as CpBINAP), 2,2'-bis(dicyclohexylphosphino)-1,1'-binaphthyl (hereinafter referred to as CyB-INAP), and 2,2'-bis(diphenylphosphino)-5,5',6,6',7,7',8, 8'-octahydro-1,1'-binaphthyl (hereinafter referred to as H₈-BINAP).

Of the above-enumerated ligands represented by formula (XVI), CHIRAPHOS and those in which A¹ is -A²-Ar-Ar-A²- or -Ar-Ar- have an asymmetric center to produce an (S)-isomer, an (R)-isomer, or a racemate, any of which can be used in the present invention. In what follows, the indication of "(S)-" or "(R)-" for the ligands having an asymmetric structure is omitted in some cases.

Of the above-described ligands, particularly preferred in the practice of the present invention are those represented by formula (XVI).

The auxiliary ligands which may be used in the ruthenium-phosphine complex include 1,5-cyclooctadiene, norbornadiene, benzene, p-cymene, mesitylene, acetonitrile, benzonitrile, pyridine, quinoline, isoquinoline, acetic acid, and acetylacetonato.

Complexes 1 to 4 represented by the following formulae (XVII) to (XX) are examples of preferred ruthenium-phosphine complexes.



wherein X¹ represents a halogen atom or R⁸COO, wherein R⁸ represents a hydrogen atom, a lower alkyl group having 1 to 4 carbon atoms or a lower haloalkyl group having 1 to 4 carbon atoms; L represents an organic phosphine compound; a and b each represent an integer of 0 to 2, giving the sum of 2; and c represents an integer of 1 to 4.



wherein X² represents ClO₄, PF₆ or BF₄; L is as defined above; where L is a monodentate ligand, e is 2, and f is 2 when d is 0, or e is 4, and f is 1 when d is 1; and where L is a bidentate ligand, e is 1, and f is 2 when d is 0, or e is 2, and f is 1 when d is 1.



wherein X³ represents a halogen atom; Bz represents a substituted or unsubstituted benzene ring; X⁴ represents a halogen atom, ClO₄, PF₆, BF₄ or BPh₄ (wherein Ph stands for a phenyl group, hereinafter the same); L is as defined above; g represents 1 or 3, where X³ and X⁴ are each an iodine atom, g may be 3; and h represents 2 where L is a monodentate ligand, or h represents 1 where L is a bidentate ligand.



wherein T represents a tertiary amine; L is as defined above; and w is 4 where L is a monodentate ligand, or w is 2

where L is a bidentate ligand.

In formulae (XVII) to (XX), the organic phosphine compound as L is selected from those enumerated above.


In formula (XVII), X^1 preferably represents a chlorine atom, a bromine atom, an iodine atom, $HCOO$, CH_2COO , and CF_3COO , with a chlorine atom being particularly preferred. The integers a and b make three combinations; ($a=0$, $b=2$), ($a=1$, $b=1$), and ($a=2$, $b=0$). The combination ($a=0$, $b=2$) is preferred. Where X^1 is a halogen atom, c is preferably 3 or 4.

In formula (XIX), the halogen atom as X^3 or X^4 includes a chlorine atom, a bromine atom, and an iodine atom. The substituted or unsubstituted benzene ring as Bz means a benzene ring which may have one or more substituents selected from an alkyl group, an alkoxy group, an alkoxycarbonyl group, a halogen atom, etc. Preferred examples of Bz include unsubstituted benzene and benzene substituted with an alkyl group having 1 to 4 carbon atoms, an alkoxy group having 1 to 4 carbon atoms, an alkoxycarbonyl group having 1 to 4 carbon atoms, a chlorine atom, a bromine atom, or an iodine atom. Specific examples are benzene, toluene, xylene, trimethylbenzene, hexamethylbenzene, ethylbenzene, *t*-butylbenzene, *p*-cymene, cumene, anisole, methyl benzoate, and chlorobenzene.


In the formula (XX), the tertiary amine as T includes triethylamine, tributylamine, ethyldiisopropylamine, 1,8-bis(dimethylamino)naphthalene, dimethylaniline, pyridine, and *N*-methylpiperidine, with triethylamine being preferred.

Preferred examples of complexes 1 to 4 are shown below.


Complex 1:




Complex 2:



Complex 3:



Complex 4:



Complexes 1 are particularly preferred among complexes 1 to 4 in view of their reaction selectivity.

The base comprising an alkali metal or an alkaline earth metal which can be used in the present invention may be either organic or inorganic, including compounds represented by formula (XXI):



wherein M represents an alkali metal or an alkaline earth metal, preferably an alkali metal; and R^9 represents a hydroxyl group, an alkoxy group having 1 to 4 carbon atoms, or a mercapto group.

Preferred examples of the base (XXI) are KOH, $Ca(OH)_2$, KOMe, KOt-Bu, LiOH, LiOMe, LiOt-Bu, NaOH, NaOMe, and $Mg(OMe)_2$ (wherein Me stands for a methyl group, and t-Bu a t-butyl group, hereinafter the same), with those having an alkali metal being still preferred. KOH and NaOH are particularly preferred.

The base is preferably used in an amount of about 0.5 to 100 equivalents, particularly about 1 to 40 equivalents, to the complex.

The amine which can be used in the present invention includes primary, secondary or tertiary amines represented by formula (XXII):



wherein R^{10} , R^{11} , and R^{12} , which may be the same or different, each represent a hydrogen atom; a substituted or unsubstituted alkyl group, a substituted or unsubstituted aralkyl group or a substituted or unsubstituted aryl group; provided that R^{10} , R^{11} , and R^{12} do not simultaneously represent a hydrogen atom; primary, secondary or tertiary diamines represented by formula (XXIII):



wherein R^{13} , R^{14} , R^{15} , and R^{16} , which may be the same or different, each represent a hydrogen atom, a substituted or unsubstituted alkyl group, a substituted or unsubstituted aralkyl group or a substituted or unsubstituted aryl group; and Z represents a substituted or unsubstituted and saturated or unsaturated hydrocarbon chain having 1 to 6 carbon atoms or a substituted or unsubstituted and saturated or unsaturated hydrocarbon ring having 3 to 6 carbon atoms; and cyclic amines.

The substituted or unsubstituted alkyl group as R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} includes straight-chain, branched or cyclic alkyl groups which may have one or more substituents, such as an alkoxy group. Straight-chain or branched alkyl groups having 1 to 10 carbon atoms and cycloalkyl groups having 5 to 8 carbon atoms are still preferred.

The substituted or unsubstituted aralkyl group as R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} includes an alkyl group substituted with an aryl group that may have one or more substituents, such as an alkyl group and an alkoxy group. The aryl moiety in the aralkyl group is preferably a substituted or unsubstituted phenyl group, and the alkyl moiety preferably has 1 to 4 carbon atoms. A benzyl group is an example.

The substituted or unsubstituted aryl group as R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , and R^{16} includes an aryl group which may have one or more substituents, such as an alkyl group and an alkoxy group. Preferred examples include a phenyl or naphthyl group which may be substituted with a straight-chain or branched alkyl group having 1 to 4 carbon atoms or a straight-chain or branched alkoxy group having 1 to 4 carbon atoms.

Specific examples of the amines include monoamines, such as methylamine, ethylamine, propylamine, isopropylamine, butylamine, isobutylamine, t-butylamine, hexylamine, octylamine, dodecylamine, cyclopentylamine, cyclohexylamine, benzylamine, dimethylamine, diethylamine, dipropylamine, diisopropylamine, dibutylamine, diisobutylamine, di-t-butylamine, dihexylamine, dicyclopentylamine, dicyclohexylamine, dibenzylamine, trimethylamine, triethylamine, tripropylamine, ethyldiisopropylamine, tributylamine, trihexylamine, tribenzylamine, benzyldimethylamine, aniline, p-toluidine, N,N-dimethylaniline, diphenylamine, triphenylamine, piperidine, piperazine, morpholine, N-methylpiperidine, N-methylpiperazine, and N-methylmorpholine; diamines, such as ethylenediamine, trimethylenediamine, tetramethylenediamine, pentamethylenediamine, hexamethylenediamine, N-methylethylenediamine, N,N'-dimethylethylenediamine, N,N'-dimethylpropylenediamine, N,N'-dimethyltetramethylenediamine, N,N'-diethylethylenediamine, N,N'-diethylpropylenediamine, N,N'-diethyltetramethylenediamine, N,N'-dibenzylethylenediamine, N,N'-dibenzylpropylenediamine, N,N'-dibenzyltetramethylenediamine, N,N'-diphenylethylenediamine, N,N'-diphenylpropylenediamine, N,N'-diphenyltetramethylenediamine, N,N,N'-trimethylethylenediamine, tetramethylethylenediamine, tetramethylpropylenediamine, tetramethyltetramethylenediamine, tetraethylethylenediamine, tetraethylpropylenedi-

amine, tetraethyltetramethylenediamine, tetrabenzyl thylenediamine, tetrabenzylpropylenediamine, tetrabenzyl-
 tramethylenediamine, tetraphenylethylenediamine, tetraphenylpropylenediamine, tetraphenyltetramethylenediamine,
 and o-phenylenediamine; and optically active diamines, such as optically active 1,2-diphenylethylenediamine, optically
 active 1,3-diphenylpropylenediamine, optically active 1,4-diphenyltetramethylenediamine, optically active 1,2-diami-
 5 nopropane, optically active 1,1-diphenyl-1,2-diaminopropane, optically active 1,1-di(p-methoxyphenyl)-1,2-diamino-
 propane, optically active 2,3-diaminobutane, optically active 2,4-diaminopentane, optically active 2,5-diaminohexane,
 optically active 1,2-diaminocyclopentane, and optically active 1,2-diaminocyclohexane.

Of these amines preferred are diamines represented by formula (XXIII). Particularly preferred are primary diamines
 (XXIII) in which R¹³, R¹⁴, R¹⁵, and R¹⁶ all represent a hydrogen atom, and Z is saturated hydrocarbon chain having 1
 10 to 4 carbon atoms, such as ethylenediamine, trimethylenediamine, and tetramethylenediamine.

The monoamine is used in an amount of about 1 to 8 equivalents, preferably about 2 to 4 equivalents, to the
 complex, and the diamine is used in an amount of about 0.5 to 4 equivalents, preferably about 1 to 4 equivalents, to
 the complex.

The process of the present invention is carried out by hydrogenating the compound (II) in the presence of the base
 15 and the amine using the ruthenium-phosphine complex as a catalyst in a hydrogen atmosphere. The catalyst is used
 in an amount of about 1/5 to 1/100000 mol, preferably about 1/200 to 1/50000 mol, per mole of the substrate, i.e., the
 compound (II) (substrate/catalyst molar ratio=about 5 to 100000, preferably about 200 to 50000). The reaction is pref-
 erably conducted with stirring. In particular, when the reaction is carried out with a small amount of the catalyst, it is
 preferable to mechanically stir the reaction system by means of a mechanical stirrer, etc.

The reaction temperature ranges usually from about -30° to 250°C, preferably from about -10° to 100°C. Within
 20 this range, a low temperature is preferred from the standpoint of reaction selectivity, and a high temperature is preferred
 for obtaining an increased reaction rate.

The reaction usually completes in several minutes to about 30 hours while varying depending on the concentration
 of the reaction substrate, the amount of the catalyst, the temperature, the hydrogen pressure, and the like. The com-
 25 pletion of the reaction is confirmed by gas chromatography, etc.

The hydrogen pressure is preferably about 1 to 200 atm., still preferably about 3 to 100 atm. Hydrogen gas can
 be used as diluted with an inert gas, such as methane, nitrogen, argon, helium, carbon dioxide, etc., either alone or
 as a mixture thereof.


The reaction can be performed with or without a solvent. Any solvent may be used with no particular limitation as
 30 long as it gives no adverse influence on the reaction. Examples of suitable solvents include water; hydrocarbons, such
 as hexane, heptane, octane, nonane, decane, benzene, toluene, and xylene; ethers, such as tetrahydrofuran, dioxane,
 dimethoxyethane, diisopropyl ether, and diethylene glycol dimethyl ether; esters, such as ethyl acetate, butyl acetate,
 ethyl propionate, and ethyl acetoacetate; alcohols, such as methanol, ethanol, n-propanol, and isopropyl alcohol; ni-
 35 triles, such as acetonitrile; phosphorous acid and esters thereof, such as phosphorous acid, trimethyl phosphite, dime-
 thyl phosphite, monomethyl phosphite, triethyl phosphite, tributyl phosphite, trioctyl phosphite, triphenyl phosphite,
 dimethylphenyl phosphite, and methyl-diphenyl phosphite; sulfoxides, such as dimethyl sulfoxide, diethyl sulfoxide,
 dibenzyl sulfoxide, diphenyl sulfoxide, and tetramethylene sulfoxide; and amides, such as formamide, N,N-dimethyl-
 formamide, N,N-dimethylacetamide, and N-methylpyrrolidone. These solvents can be used either individually or as a
 combination of two or more thereof. Of these solvents preferred are alcohols, such as methanol, ethanol, and isopropyl
 40 alcohol. Isopropyl alcohol is a particularly preferred solvent.

While the amount of the solvent, if used, is not particularly limited, it is preferably about 0.5 to 100 times the weight
 of the reaction substrate.

After completion of the reaction, the reaction mixture is purified in a conventional manner by, for example, filtration,
 concentration under reduced pressure, distillation, and the like.

The threshold value of the odor of the compound (I) was determined by a triangle method well-known in the art
 45 (see T.E. Acree, et al., *J. Agric. Food Chem.*, Vol. 33, pp. 425-427 (1985)). It was revealed as a result that the threshold
 value of (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (IX) is 1/40 of that of its enantiomer, i.e.,
 (E)-(S)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (XI) and that the threshold value of (E)-(R)-2-me-
 thyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (XXIV) is 1/8 of that of its enantiomer, i.e., (E)-(S)-2-methyl-4-
 50 (2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (XXV). In other words, the odor of the (E-R)-ethyl compound (IX) is
 about 40 times as strong as the (E-S)-ethyl compound (XI), and the odor of the (E-R)-methyl compound (XXIV) is about
 8 times as strong as its enantiomer (XXV).


The (E-R)-ethyl compound (IX) has a clean and warm sandalwood oil odor, while the (Z-R)-ethyl compound (XXVI)
 is inferior in strength of odor to the compound (IX), and its odor has butter-like sweetness, lacking a woody note. Thus,
 55 the (E-R)-ethyl compound (IX) has an excellent sandalwood oil odor, and responsible for the odor characteristic of the
 compound according to the present invention.




(IX)




(XXIV)



(XI)



(XXV)



(XXVI)

In carrying out the present invention, the starting (1S,5S)- α -pinene does not always need to be optically pure. The inventors prepared an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol having an optical purity of about 50%e.e. starting from (1S,5S)- α -pinene having an optical purity of about 50%e.e. that is available with relative ease. Substantially indistinguishable in odor from a substantially optically pure compound, the resulting compound achieves the object of the present invention and is economically more advantageous.

It has been thus confirmed that the compound (I) has an excellent sandalwood oil odor that is more pleasant and stronger than the odor of its enantiomer, (E)-(S)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol, or a racemic mixture of the compound (I) and its enantiomer.

Blending the compound (I) provides perfume compositions having a fresh and appealing sandalwood oil odor and serving as an odorant, a modifier, an intensifier, and the like. The content of the compound (I) in the perfume composition of the invention is not particularly limited and is decided appropriately in accordance with the purpose. For example, the compound (I) is usually used in an amount of about 0.1 to 50% by weight based on the total composition. If desired, the perfume composition can contain compounded perfumes or bases for compounding commonly employed in the art.

The perfume composition of the invention is useful in various cosmetics, fragrances, toiletries, and the like. More specifically, the perfume composition is added to toiletries, such as soaps, shampoos, rinses, tooth pastes, and mouth washes; hair care cosmetics, such as hair dyes and hair tonics; cosmetics for foundation, such as creams, lotions, and face packs; cosmetics for make-up, such as face powder, foundation, and rouge; fragrances, such as perfumes and eau-de-cologne; suntan lotions or sunscreens; lip cosmetics, such as lip sticks and lip creams; deodorants and room odorant; hygienic goods, such as disinfectants and insecticides; bleach, softeners, dishwashing detergents, and so on, in an appropriate amount for adding its odor to thereby increase their commercial value.

As described above, the present invention provides a novel optically active compound, (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol, in high yield with ease. This compound has a strong and excellent sandalwood oil odor clearly distinguishable from the odor of conventional synthetic compounds, and the perfume compositions containing the same are effectively used in wide fields including cosmetics, fragrances, toiletries, and hygienic goods.

The present invention will now be illustrated in greater detail with respect to Examples, but it should be understood that the present invention is not construed as being limited thereto.

In Examples, various analyses were made with the following instruments.

Gas Chromatography (GC):

Gas chromatograph: HP-5890, manufactured by Hewlett Packard
 Column: PEG BC-WAX (0.25 mm x 50 mm), manufactured by GL-Science K.K.
 Measuring temperature: 100 to 220°C (increased at a rate of 10°C/min)
 Injection temperature: 250°C
 Retention time:

EP 0 829 463 A2

- (1) (E)-(R)-2-Alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol: about 12.7 min (area intensity: a_1)
(2) Other isomers: about 11.2 to 12.4 min (area intensity: a_2)
(3) (E)-(R)-2-Alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al: about 10.3 min (area intensity: a_3)
(4) (E)-(S)-2-Alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al: about 9.4 min (area intensity: a_4)

Conversion (%): $[(a_1+a_2)/(a_1+a_2+a_3+a_4)] \times 100$

Reaction selectivity (%): $[a_1/(a_1+a_2)] \times 100$

Preparative High-Performance Liquid Chromatography (HPLC):

Chromatograph: PU-615, manufactured by GL-Science K.K.

Column: INERTSIL ODS-3 (20 x 250 mm), manufactured by Nakarai Tesc K.K.

Solvent: methanol

Flow rate: 10 ml/min

Detector: RI-98 (refractive index: preparative), manufactured by Labo System

Infrared Absorption Spectrum (IR):

IR spectrophotometer: IR-810, JASCO Inc.

Measuring method: liquid film method

Mass Spectrum (MS):

Mass spectrometer: M-80B (ionization voltage: 20 eV), manufactured by Hitachi, Ltd.

Proton Nuclear Magnetic Resonance ($^1\text{H-NMR}$):

NMR spectrometer: AM-400 (400 MHz), manufactured by Bruker, Inc.

Internal standard: tetramethylsilane

Specific Rotation:

Polarimeter: DIP-370, manufactured by JASCO Inc.

EXAMPLE 1

Synthesis of (E)-(R)-2-Methyl-4-(2, 2,3-trimethylcyclopent-3-en-1-yl)-2-butene-1-ol:

In a 100 ml stainless steel autoclave were charged 3.84 g (0.02 mol) of (E)-(R)-2-methyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al synthesized from (1S,5S)- α -pinene having an optical purity of 97% e.e. (available from Aldrich) in a conventional manner, 38.4 mg (0.04 mmol) of $\text{RuCl}_2(\text{PPh}_3)_3$, 4 ml of a 0.02M 1,3-diaminopropane (0.08 mmol) solution in isopropyl alcohol, and 12 ml of a 0.1M potassium hydroxide (1.2 mmol) solution in isopropyl alcohol in a nitrogen atmosphere, and the mixture was stirred at 0°C for 3 hours under a hydrogen pressure of 50 atm.

The analysis of the reaction mixture by GC revealed that the conversion was 100% and the reaction selectivity was 98%. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure. The residue was dissolved in hexane, washed with a saturated sodium chloride aqueous solution, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting oily substance was distilled under reduced pressure to give 3.13 g (81%) of the title compound. The chemical purity of the product was found to be 98% by GC.

Boiling point: 125.7 to 126°C (5 mmHg)

Specific rotation: $\alpha_D^{25} = +0.71^\circ$

MS (m/z): 194(M⁺), 179, 161, 135, 121, 109, 95, 81, 67

IR (film) (cm⁻¹): 3325, 2950, 1460, 1440, 1360, 1015

$^1\text{H-NMR}$ (CDCl_3 , δ ppm): 5.44 (1H, t, J=7.4Hz), 5.22 (1H, brs), 4.01 (2H, brs), 2.28-1.99 (3H, m), 1.84-1.74 (2H, m), 1.69 (3H, s), 1.60 (3H, s), 1.45 (1H, s), 0.99 (3H, s), 0.80 (3H, s)

(E)-(S)-2-Methyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al synthesized from (1R,5R)- α -pinene having an optical purity of 97%e.e. (available from Aldrich) was hydrogenated in the same manner as in Example 1 to obtain 3.08 g (yield 80%) of the title compound. The chemical purity of the product was found to be 98% by GC. Specific rotation: $\alpha_D^{25} = -0.71^\circ$

The boiling point, MS, IR, and ^1H -NMR data of the product were the same as those of Example 1.

Synthesis of (E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:

In a 100 ml stainless steel autoclave were charged 4.12 g (0.02 mol) of (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al synthesized from (1S,5S)- α -pinene having an optical purity of 97% e.e. in a conventional manner, 38.4 mg (0.04 mmol) of $\text{RuCl}_2(\text{PPh}_3)_3$, 4 ml of a 0.02M 1,3-diaminopropane (0.08 mmol) solution in isopropyl alcohol, and 12 ml of a 0.1M potassium hydroxide (1.2 mmol) solution in isopropyl alcohol in a nitrogen atmosphere, and the mixture was stirred at 0°C for 3 hours under a hydrogen pressure of 50 atm.

The analysis of the reaction mixture by GC revealed that the conversion was 100% and the reaction selectivity was 98%. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure. The residue was dissolved in hexane, washed with a saturated sodium chloride aqueous solution, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting oily substance was distilled under reduced pressure to give 3.55 g (yield 85%) of (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol. As a result of GC, the product had a chemical purity of 95% and contained the (Z)-compound in a proportion of 2%.

Boiling point: 140 to 140.5°C (12 mmHg)
 Specific rotation: $\alpha_D^{25} = -2.9^\circ$
 MS (m/z): 208(M⁺), 193, 175, 161, 145, 135, 121, 109, 108, 95, 93, 79, 67
 IR (film) (cm⁻¹): 3325, 2950, 1460, 1440, 1360, 1015
¹H-NMR (CDCl₃, δ ppm): 5.41 (1H, t, J=7.1Hz, 3-CH=C), 5.22 (1H, brs), 4.05 (2H, brs, 1-CH₂), 2.28-2.17 (2H, m), 2.14 (2H, q, J=7.5Hz, 5-CH₂), 2.04-1.95 (1H, m), 1.85-1.77 (2H, m), 1.61 (3H, s), 1.30 (1H, brs), 1.01 (3H, t, J=7.5Hz, 6-Me), 1.00 (3H, s), 0.81 (3H, s)

The ^1H - ^1H -Cosy and ^1H - ^1H -NOESY spectra displayed a nucleo-overhauser effect (NOE), providing confirmation of the (E)-configuration.

The resulting (R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol (1.0 g) having an (E)/(Z) ratio of 98/2 was dissolved in 2.0 ml of methanol and purified by recycle operation of a preparative HPLC. After recycling, 70 mg of a fraction having an (E)/(Z) ratio of about 1/1 was collected, which was dissolved in 1.2 ml of methanol, and again subjected to recycle operation to obtain the (Z)-compound and the (E)-compound in a yield of 4 mg each.

¹H-NMR (CDCl₃, δ ppm): 5.34 (1H, dd, J=7.1 and 7.7Hz, 3-CH=C), 5.22 (1H, brs), 4.18 (2H, d, J=7.4Hz, 1-CH₂), 2.0-2.3 (5H, m), 1.7-1.9 (2H, m), 1.60 (3H, m), 1.08 (1H, t, J=5.6Hz), 1.04 (3H, t, J=7.4Hz, 6-Me), 1.00 (3H, s), 0.80 (3H, s)

The ^1H - ^1H -Cosy and ^1H - ^1H -NOESY spectra displayed the NOE effect, providing confirmation of the (Z)-configuration.

The NMR spectrum of the (E)-compound was the same as the product before purification.

EXAMPLES 3 TO 7Synthesis of (E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:

In the same manner as in Example 2, the title compound was prepared by hydrogenating (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al which was synthesized from (1S,5S)- α -pinene having an optical purity varying from 50%e.e. to 92%e.e.

The boiling points, MS, IR, and ^1H -NMR data of all the products obtained were the same as those in Example 2. The optical purity and specific rotation of the products are shown in Table 1 below.

TABLE 1

	Example No.				
	3.	4	5	6	7
Optical Purity (%e.e.)	92	80	70	60	50
Specific rotation ($^\circ$)	-2.7	-2.4	-2.1	-1.8	-1.5

REFERENCE EXAMPLE 2Synthesis of (E)-(S)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:

(E)-(S)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-al synthesized from (1R,5R)- α -pinene having an optical purity of 97%e.e. (available from Aldrich) was hydrogenated in the same manner as in Example 2 to obtain the title compound.

Specific rotation: $\alpha_D^{25} = +2.9^\circ$

The boiling point, MS, IR, and ^1H -NMR data were the same as those of Example 2.

EXAMPLE 8Synthesis of (E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:

Hydrogenation reaction was carried out in the same manner as in Example 2, except that the reaction was carried out under a hydrogen pressure of 50 atm. at room temperature for 2 hours while stirring.

The results of GC revealed that the conversion was 100%, and the reaction selectivity was 94%.

As is apparent from the comparison between Example 2 and Example 8, the reaction sensitivity is higher at 0°C than at room temperature.

EXAMPLES 9 TO 15Synthesis of (E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:

Hydrogenation reaction was conducted in the same manner as in Example 2, except for changing the kinds and amounts of the diamine and the base. The reaction results are shown in Table 2 below.

TABLE 2

Example No.	Diamine		Base		Conversion (%)	Selectivity (%)
	Kind	Amount (ml*)	Kind	Amount (ml**)		
9	1,2-diaminoethane	4	KOH	12	99.1	93.6
10	1,3-diaminopropane	4	KOH	20	99.4	94.0
11	1,3-diaminopropane	4	NaOH	12	99.2	94.3

Note: * 0.02M Solution in isopropyl alcohol.

** 0.2M Solution in isopropyl alcohol.

TABLE 2 (continued)

Example No.	Diamine		Base		Conversion (%)	Selectivity (%)
	Kind	Amount (ml*)	Kind	Amount (ml**)		
12	1,3-diamino-propane	4	t-BuOK	12	99.0	91.1
13	1,3-diamino-propane	4	MeONa	12	99.1	89.4
14	1,3-diamino-propane	8	KOH	12	99.2	94.2
15	1,3-diamino-propane	16	KOH	12	99.3	95.8

Note: * 0.02M Solution in isopropyl alcohol.

** 0.2M Solution in isopropyl alcohol.

EXAMPLES 16 TO 23**Synthesis of (E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:**

Hydrogenation reaction was conducted in the same manner as in Example 2, except for changing the kind and amount of the catalyst and the kind of the diamine and using a solvent as shown in Table 3 below. The reaction results are shown in Table 3.

TABLE 3

Example No.	Catalyst		Diamine	Solvent	Conversion (%)	Selectivity (%)
	Kind	Amount (mg)				
16	RuCl ₂ (Ph ₃ P) ₃	38.4	1,3-diamino-propane	toluene	99.0	93.6
17	RuCl ₂ (Ph ₃ P) ₃	38.4	1,3-diamino-propane	diethyl ether	99.0	94.0
18	RuCl ₂ (Ph ₃ P) ₃	38.4	1,3-diamino-propane	MeCN	97.1	91.3
19	RuCl ₂ (Ph ₃ P) ₃	38.4	1,3-diamino-propane	hexane	99.1	94.2
20	RuCl ₂ (Ph ₃ P) ₃	38.4	1,3-diamino-propane	tetra-hydrofuran	99.0	92.4
21	RuCl ₂ (Ph ₃ P) ₃	16.9	1,2-diamino-ethane	isopropyl alcohol	99.7	96.3
22	RuCl ₂ (Ph ₃ P) ₃	8.4	1,2-diamino-ethane	isopropyl alcohol	99.6	95.4
23	Ru-BINAP catalyst*	38.4	1,3-diamino-propane	isopropyl alcohol	97.5	85.5

Note: * [Ru₂Cl₄(S)-To-BINAP]₂(NEt)₃**EXAMPLE 24****Synthesis of (E)-(R)-2-Propyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:**

In a 100 ml stainless steel autoclave were charged 6.60 g (0.03 mol) of (E)-(R)-2-propyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol synthesized from (1S)- α -pinene having an optical purity of 97% in a conventional

EP 0 829 463 A2

manner, 28.7 mg (0.03 mmol) of $\text{RuCl}_2(\text{PPh}_3)_3$, 2.4 ml of a 0.1M 1,2-diaminoethane (0.24 mmol) solution in isopropyl alcohol, and 15 ml of a 0.1M potassium hydroxide (1.5 mmol) solution in isopropyl alcohol in a nitrogen atmosphere, and the mixture was stirred at room temperature for 3 hours under a hydrogen pressure of 50 atm.

The analysis of the reaction mixture by GC revealed that the conversion was 100%. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure. The residue was dissolved in hexane, washed with a saturated sodium chloride aqueous solution, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The resulting oily substance was distilled under reduced pressure to give 4.80 g (72%) of the title compound.







Boiling point: 137.5 to 138°C (4 mmHg)
Specific rotation: $\alpha_D^{25} = -0.23^\circ$
MS (m/z): 222(M⁺), 207, 189, 161, 147, 135, 121, 108, 95, 81, 71
IR (film) (cm⁻¹): 3320, 2950, 1460, 1380, 1360, 1005
¹H-NMR (CDCl₃, δ ppm): 5.46 (1H, t, J=7.2Hz), 5.22 (1H, brs), 4.02 (2H, brs), 2.16-2.29 (2H, m), 2.10 (2H, dd, J=7.2, 8.4Hz), 1.99-2.01 (1H, m), 1.76-1.85 (2H, m), 1.60 (4H, brs), 1.44 (2H, q, J=7.6Hz), 1.00 (3H, s), 0.92 (3H, t, J=7.2Hz), 0.80 (3H, s)

EXAMPLE 25

Evaluation of Quality of Odor:

The odor of the compounds prepared in Examples 1, 2 and 24 and Reference Examples 1 and 2 was evaluated by 7 specialized panelists in accordance with a triangle method (see T. E. Acree, et al., J. Agric. Food Chem., Vol. 33, pp. 425-427 (1985)). The average of the threshold values given by 7 panelists is shown in Table 4 below.

TABLE 4

Preparation Example No.	Configuration	Compound	Evaluation on Quality of Odor	Threshold Value
Example 1	(E)-(R)		fresh and strong sandalwood oil odor associated with green trees	51
Reference Example 1	(E)-(S)		dry and weak sandalwood oil odor with a milky and floral note	424
Example 2	(E)-(R)		clean, bright and strong sandalwood oil odor with richness and a woody note	0.85
Example 2	(Z)-(R)		inferior to the trans-compound in strength, lacking a woody note, and having butter-like sweetness	unmeasured
Reference Example 2	(E)-(S)		milky sandalwood oil odor with a cedar character	44
Example 24	(E)-(R)		soft and warm sandalwood oil odor, slightly weaker than the (E)-(R)-compound of Example 2	unmeasured

Note: * 10⁻¹² M/lEXAMPLE 26Evaluation on Quality of Odor vs. Optical Purity:

The quality of odor of the compounds obtained in Examples 2 to 7 was evaluated by 7 panelists. The results obtained are shown in Table 5 below.

COMPARATIVE EXAMPLE 1

(E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol having an optical purity of 40%e.e. was synthesized from (1R,5R)- α -pinene having an optical purity of 40%e.e. in the same manner as in Example 2. The odor of the resulting compound was evaluated by 7 panelists. The results obtained are shown in Table 5.

COMPARATIVE EXAMPLE 2

The odor of commercially available 2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol ((R)/(S)=30/70), Baccanol (registered trademark, available from IFF) was evaluated by 7 panelists. The results obtained are shown in Table 5.

TABLE 5

Example No.	Optical Purity (%e.e.)	Evaluation on Quality of Odor	Diffusibility	Volume and Depth
Example 2	97	strong, clear, warm and creamy odor primarily having a woody note inherent to natural sandalwood	present	present
Example 3	92	strong, clear, warm and creamy odor primarily having a woody note inherent to natural sandalwood	present	present
Example 4	80	strong, clear, warm and creamy odor primarily having a woody note inherent to natural sandalwood	present	present
Example 5	70	strong odor having a woody note inherent to natural sandalwood, slightly associated with green trees	present	present
Example 6	60	strong odor having a woody note inherent to natural sandalwood, slightly associated with green trees	present	present
Example 7	50	sandalwood oil odor, while slightly astringent compared with Examples 2 to 6	present	present
Comparative Example 1	40	sandalwood oil odor as in Examples 2 to 7 but lacking clearness with increased cedarwood-like astringency and sweetness	absent	absent
Comparative Example 2	Bacdanol	less sandalwood-like than Examples 2 to 7	absent	absent

EXAMPLE 27**Floral Green and Musky Perfume Composition for Toiletries:**

A floral green and musky perfume composition for toiletries was prepared using the (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol prepared in Example 3 in accordance with the following formulation (unit: part by weight, hereinafter the same).

	Abbalide 50 BB (registered trademark) ¹⁾	50
10	Bergamot oil	50
	Cyclamenaldehyde ²⁾	80
	Eugenol	2
	γ -Methylionone ³⁾	30
15	Geraniol	10
	Methyl dihydrojasmonate	100
	Helional ⁴⁾	80
	Hexylcinnamaldehyde	30
	Phenylacetaldehyde	2
20	Kovanol (registered trademark) ⁵⁾	100
	1-Citronellol	60
	1-Citronelyl acetate	3
	1-Hydroxycitroellal	20
25	Lilial ⁶⁾	90
	Linalool	30
	Linalyl acetate	6
	Ethylene brassylate	150
	Floralozone (registered trademark) ⁷⁾	3
30	Allylamyl glycolate	1
	2-Phenylethanol	33
	(E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol obtained in Example 3	70
	Total:	1000

1) A 50% solution of 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta(γ)-2-benzopyran in benzyl benzoate (available from Bush Boak Allen)

2) p-Isopropyl- α -methylhydrocinnamaldehyde

3) 4-(2,2,6-Trimethyl-2-cyclohexen-1-yl)-3-methyl-3-buten-3-one

4) α -Methyl-3,4-methylenedioxyhydrocinnamaldehyde

5) 4(3)-(4-Hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxyaldehyde (available from Takasago International Corporation)

6) p-t-Butyl- α -methylhydrocinnamaldehyde

7) p-Ethyl- α , α -dimethylhydrocinnamaldehyde (available from IFF)

EXAMPLE 28**Floral Perfume Composition for Soaps:**

A highly appealing floral perfume composition for soaps was prepared using the (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol prepared in Example 4 in accordance with the following formulation.

	Abbalide 50 BB (registered trademark)	110
50	Decyl aldehyde	2
	Undecyl aldehyde	6
	Benzyl salicylate	30
	Linalool	50
55	1-Citronellol	130
	1-Citronelylnitrile	2
	Tricyclodecenyl propionat	10

(continued)

	α -Damascone	1
	Ethyl vanillin	3
5	Hexylcinnamaldehyde	140
	Indole ¹⁾	10
	Iso E Super (registered trademark) ²⁾	100
	Coumarin	20
10	Heliopropine	5
	Benzyl acetate	80
	Lilial	50
	γ -Methylionone	40
	Orange oil (PERA BRAZIL)	20
15	2-Phenylethanol	90
	Rose oxide	5
	9-Decenol	1
	Rosephenone	10
0	p-t-Butylcyclohexyl acetate	40
	(E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol obtained in Example 4	45
	Total:	1000

1) 10% Dipropylene glycol solution

2) 7(6)-Acetyl-1.2.3.4.5.6.7.8-octahydro-1.1.6.7-tetraethylnaphthalene (available from IFF)

EXAMPLE 29**Floral Perfume Composition for Shampoos:**

A highly appealing floral perfume composition for shampoos was prepared using the (E)-(R)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol prepared in Example 4 in accordance with the following formulation:

	γ -Undecalactone ¹⁾	15
35	Ambroxan (registered trademark) ^{2,3)}	1
	Benzyl acetate	50
	Bergamot oil	50
	Citronellol	40
40	Ethyl (1S,6R)-2,2,6-trimethylcyclohexanecarboxylate	5
	Ethyl acetoacetate ³⁾	10
	Ethyl vanillin	2
	Eugenol	15
	γ -Methylionone	17
45	Methyl dihydrojasmonate	30
	Helional (registered trademark) ⁴⁾	8
	Heliotropin	20
	Hexylcinnamaldehyde	140
	Iso E Super (registered trademark)	30
50	Kovanol (registered trademark)	50
	Lemon oil (produced in California)	200
	cis-3-Hexenyl acetate	5
	Allylamyl glycolate	5

1) 1% Dipropylene glycol solution

2) Decahydro-3a.6.6.9a-tetramethylnaphtho[1.2-b]furan (available from Henkel)

3) 10% Dipropylene glycol solution

4) α -Methyl-3,4-methylenedioxyhydrocinnamaldehyde (available from IFF)

(continued)

Linalool	40
Methyl anthranilate	10
Ethylene brassylate	10
n-Hexyl salicylate	50
Oak Moss No. 1 ⁵⁾	5
p-Cresyl methyl ether	2
2-Phenylethanol	70
Tonalid (registered trademark) ⁶⁾	50
2,4-Di-t-butylcyclohexanone	20
(E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol obtained in Example 4	50
Total:	1000

5) Methyl 2,4-dihydroxy-3,6-dimethylbenzoate (produced by Takasago International Corporation)

6) 6-Acetyl-1,1,2,4,4,7-hexamethyltetralin (available from Polaks Eurutal Workobv).

COMPARATIVE EXAMPLE 3

A floral perfume composition was prepared using (E)-(R)-2-ethyl-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol obtained in Example 2 or, for comparison, Bacdanol (registered trademark; 2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol; (R)/(S) = 30/70) according to the following formulation.

Bergamot oil	20
Linalool	100
Benzyl acetate	70
Phenylacetaldehyde ¹⁾	50
Dimethylbenzylcarbinol	20
1-Hydroxycitronellal	40
1-Citronellol	80
Geraniol	50
Kovanol (registered trademark)	145
H lional	70
Lilial	90
Cyclamenaldehyde	70
Hexylcinnamaldehyde	70
2-Phenylethanol	10
Methyl dihydrojasmonate	20
Eugenol	10
Indol ¹⁾	20
Ethylene brassylate	15
(E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol obtained in Example 2 or Bacdanol	50
Total:	1000

Note: 1) 10% benzyl benzoate solution

The quality of the odor of the two perfume compositions was evaluated by 8 specialized panelists in accordance with a triangle method (one sample of the composition containing the (E-R)-compound and two samples of the composition containing Bacdanol were presented to panelists given with no preliminary information, and each panelist chose a sample having the strongest sandalwood-like odor). Six out of 8 panelist chose the composition containing the (E-R)-compound.

REFERENCE EXAMPLE 3

Synthesis of (R)-MTPA Ester of (E)-(R)-2-Ethyl-4-(2,2,3-Trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:

To 31.5 mg (151 μ mol) of (E)-(R)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol obtained in Example 2 were added 54.8 mg (234 μ mol) of (R)-MTPA-Cl, 103.5 mg (502 μ mol) of dicyclohexylcarbodiimide, and 1.0 g (12.6 mmol) of pyridine, and the mixture was stirred at room temperature for 20 hours. The reaction mixture was quenched in a 5% hydrochloric acid aqueous solution. The organic layer was extracted with hexane, and the released urea derivative was separated by filtration. The filtrate was washed successively with a 5% hydrochloric acid aqueous solution, a saturated sodium hydrogencarbonate aqueous solution, and a saturated sodium chloride aqueous solution, dried over anhydrous sodium sulfate, followed by filtration. The solvent was removed by evaporation, and the residue was purified by silica gel column chromatography using toluene as a solvent to obtain the title compound.

REFERENCE EXAMPLE 4


Synthesis of (R)-MTPA Ester of (E)-(S)-2-Ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol:

An (R)-MTPA ester of (E)-(S)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol was prepared from the (E)-(S)-2-ethyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol obtained in Reference Example 2 in the same manner as in Reference Example 3.

While the invention has been described in detail and with reference to specific examples thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

Claims


1. An (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol represented by formula (I):



(I)


wherein R represents a straight-chain or branched alkyl group having 1 to 3 carbon atoms.

2. A compound as claimed in Claim 1 having an optical purity of 50%e.e. or higher.
3. A process for preparing an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol represented by formula (I):



(I)


wherein R represents a straight-chain or branched alkyl group having 1 to 3 carbon atoms, comprising hydrogenating an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol represented by formula (I):



(II)

wherein R is as defined above,
in the presence of a ruthenium-phosphine complex as a catalyst, a base comprising an alkali metal or an alkaline earth metal, and an amine.


4. A perfume composition containing an (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol represented by formula (I):



(I)

wherein R represents a straight-chain or branched alkyl group having 1 to 3 carbon atoms.

5. A perfume composition as claimed in Claim 4 wherein the compound represented by formula (I) has an optical purity of 50%e.e. or higher.



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 829 463 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:

02.06.1999 Bulletin 1999/22

(51) Int Cl.⁶: C07C 33/12, C11B 9/00

(43) Date of publication A2:

18.03.1998 Bulletin 1998/12

(21) Application number: 97307074.1

(22) Date of filing: 11.09.1997

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

Designated Extension States:

AL LT LV RO SI

(30) Priority: 17.09.1996 JP 244842/96

16.04.1997 JP 98760/97

(71) Applicant: Takasago International Corporation
Tokyo 108 (JP)

(72) Inventors:

- Aida, Takashi, c/o Takasago Int. Corp.
Hiratsuka-shi, Kanagawa (JP)

- Harada, Makoto, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

- Iwai, Hisao, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

- Amano, Akira, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

- Yamasaki, Tetsuro, c/o Takasago Int. Corp.

Hiratsuka-shi, Kanagawa (JP)

- Yamamoto, Takeshi

Minato-ku, Tokyo (JP)

(74) Representative: Bubbs, Antony John Allen et al
GEE & CO.

Chancery House

Chancery Lane

London WC2A 1QU (GB)

(54) (E)-(R)-2-Alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol, process for preparing the same, and use thereof in perfume compositions

(57) An (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol represented by formula (I):



(I)

wherein R represents an alkyl group having 1 to 3 carbon atoms,
and a process for preparing the same comprising hydrogenating a corresponding (E)-(R)-2-alkyl-4-(2,2,3-trimethylcyclopent-3-en-1-yl)-2-buten-1-ol in the presence of a ruthenium-phosphine complex as a catalyst, a base comprising an alkali metal or an alkaline earth metal, and an amine. The compound (I) has an excellent sandalwood oil odor.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 97 30 7074

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D,X	REINHARDT R ET AL: "Enantiomer separation of alpha-campholene and fencholene derivatives by capillary gas chromatography on permethylated cyclodextrin phases I. Compounds separable with single columns" JOURNAL OF CHROMATOGRAPHY A, vol. 697, no. 1, 21 April 1995, page 475-484 XP004023082 Amsterdam, NL * compound 27 *	1-5	C07C33/12 C11B9/00
D,A	PATENT ABSTRACTS OF JAPAN vol. 005, no. 012 (C-040), 24 January 1981 & JP 55 139330 A (TAIYO KORYO), 31 October 1980 * abstract *	1,4	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07C C11B
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12 April 1999	Examiner English, R
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.92 (P+Cl.1)